

Purdue University
Purdue e-Pubs

LARS Symposia

Laboratory for Applications of Remote Sensing

1-1-1980

The Maximum Likelihood Method for Estimating Argentine Crop and Soil Test Sites Using Remote Sensing Data

Juana M. Cardoso

Mirta Aida Raed

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Cardoso, Juana M. and Raed, Mirta Aida, "The Maximum Likelihood Method for Estimating Argentine Crop and Soil Test Sites Using Remote Sensing Data" (1980). *LARS Symposia*. Paper 393.
http://docs.lib.purdue.edu/lars_symp/393

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

and

Soil Information Systems

and

Remote Sensing and Soil Survey

June 3-6, 1980

Proceedings

The Laboratory for Applications of Remote Sensing

Purdue University
West Lafayette
Indiana 47907 USA

IEEE Catalog No.
80CH1533-9 MPRSD

Copyright © 1980 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

THE MAXIMUM LIKELIHOOD METHOD FOR ESTIMATING ARGENTINE CROP AND SOIL TEST SITES USING REMOTE SENSING DATA

DR. JUANA MARIA CARDOSO
LIC. MIRTA AIDA RAED

C.N.I.E. Argentina

Our object is to study Pattern Recognition of different kind of crops in Argentine training areas by means of Maximum Likelihood Estimators just to get their multispectral signatures or distinctive spectral characteristics.

We suppose multivariate normal populations with means μ_k , covariance matrix A^k and normal density functions:

$$f(X) = (2\pi)^{-n/2} |A^k|^{-1/2} \exp \left[-\frac{1}{2} (X - \mu^k)^T (A^k)^{-1} (X - \mu^k) \right]$$

where n is the dimension of X .

Given a random sample X_1, \dots, X_n of size n , the likelihood function is given by

$$L(\theta_1, \theta_2, \dots, \theta_k) = \prod_{i=1}^n f(X_i, \theta_1, \theta_2, \dots, \theta_k)$$

The i^{th} pattern will be treated as a column vector X_i of high dimensionality, that measures the spectral reflectance of a pixel of land and assigns statistically that plot to a class of land of use.

$$X = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{pmatrix}$$

Besides we'll try to study the application of the maximum likelihood principle to pattern classification making use of Bayes' formula for conditional probabilities.